

## Advil Tablets, Caplets, Gel Caplets



### Directions

Adults and children over 12: Take 1 or 2 tablets, caplets or gel caplets every 4 hours as needed. Do not exceed 6 tablets, caplets or gel caplets in 24 hours, unless directed by a physician. Children 12 and under should use Children's Advil.

### • Caution

Keep out of reach of children. This package contains enough medicine to seriously harm a child.

### Warning

Do not take Advil if taking acetylsalicylic acid (ASA) or other products containing ibuprofen, or if allergic to ASA, salicylates or anti-inflammatory drugs, or any of its ingredients. Consult your physician before taking Advil if you have peptic ulcers, high blood pressure, heart, kidney or liver disease, any other serious disease, or are taking any other drug. Do not take Advil if you are pregnant or nursing a baby, unless directed by a physician. Consult your physician if the pain or fever persists for more than 5 days. Do not exceed the recommended dose unless advised by a physician.

If abdominal pain, heartburn, nausea or vomiting, bloating, diarrhea or constipation, ringing or buzzing in the ears, nervousness, sleeplessness, dizziness, any change in vision, fluid retention, itching, skin rashes or any other side effect or unexplained symptom develops while taking Advil, discontinue use immediately and contact a physician. In case of overdose, call a Poison Control Centre or a doctor at once even if there are no symptoms.

### Tablets and Caplets Non-medicinal Ingredients

acetylated monoglycerides, beeswax, carnauba wax, corn starch, croscarmellose sodium, ethoxyethanol, iron oxides, lecithin, microcrystalline cellulose, parabens, pharmaceutical glaze, pharmaceutical shellac, povidone, pregelatinized starch, silicon dioxide, simethicone, sodium benzoate, sodium lauryl sulfate, stearic acid, sucrose, titanium dioxide.

### Gel Caplets Non-medicinal Ingredients

corn starch, croscarmellose sodium, FD&C red No. 40, FD&C yellow No. 6, gelatin, glycerin, hypromellose, iron oxides, medium chain triglycerides, pregelatinized starch, propyl gallate, propylene glycol, silicon dioxide, sodium lauryl sulfate, stearic acid, titanium dioxide, triacetin.

## Advil Extra Strength Caplets



### Directions

Adults and children over 12: Take 1 caplet every 4 hours as needed. Do not exceed 3 caplets in 24 hours, unless directed by a physician. Children 12 and under should use Children's Advil.

### • Caution

Keep out of reach of children. This package contains enough medicine to seriously harm a child.

### Warning

Do not take Advil if taking acetylsalicylic acid (ASA) or other products containing ibuprofen, or if allergic to ASA, salicylates or anti-inflammatory drugs, or any of its ingredients. Consult your physician before taking Advil if you have peptic ulcers, high blood pressure, heart, kidney or liver disease, any other serious disease, or are taking any other drug. Do not take Advil if you are pregnant or nursing a baby, unless directed by a physician. Consult your physician if the pain or fever persists for more than 5 days. Do not exceed the recommended dose unless advised by a physician.

If abdominal pain, heartburn, nausea or vomiting, bloating, diarrhea or constipation, ringing or buzzing in the ears, nervousness, sleeplessness, dizziness, any change in vision, fluid retention, itching, skin rashes or any other side effect or unexplained symptom develops while taking Advil, discontinue use immediately and contact a physician. In case of overdose, call a Poison Control Centre or a doctor at once even if there are no symptoms.

### Non-medicinal Ingredients

corn starch, croscarmellose sodium, hydroxypropyl methylcellulose, iron oxides, lecithin, pharmaceutical shellac, polyethylene glycol, pregelatinized starch, silicon dioxide, simethicone, sodium lauryl sulfate, stearic acid, talc, titanium dioxide.

## Advil Liqui-Gels® and Extra Strength Liqui-Gels®



### Advil Liqui-Gels (200 mg) Directions

Adults and children over 12: For migraine headaches, take 1 or 2 Liqui-Gels at the first sign of symptoms and every 4 hours as needed. For all other uses, take 1 or 2 Liqui-Gels every 4 hours as needed. Do not exceed 6 Liqui-Gels in 24 hours, unless directed by a physician. Children 12 and under should use Children's Advil.

### Advil Extra Strength Liqui-Gels (400 mg) Directions

Adults and children over 12: For migraine headaches, take 1 Extra Strength Liqui-Gel at the first sign of symptoms and every 4 hours as needed. For all other uses, take 1 Extra Strength Liqui-Gel every 4 hours as needed. Do not exceed 3 Extra Strength Liqui-Gels in 24 hours, unless directed by a physician. Children 12 and under should use Children's Advil.

### • Caution

Keep out of reach of children. This package contains enough medicine to seriously harm a child.

### Warning

Do not take Advil if taking acetylsalicylic acid (ASA) or other products containing ibuprofen, or if allergic to ASA, salicylates or anti-inflammatory drugs, or any of its ingredients. Consult your physician before taking Advil if you have peptic ulcers, high blood pressure, heart, kidney or liver disease, any other serious disease, or are taking any other drug. Do not take Advil if you are pregnant or nursing a baby, unless directed by a physician. Consult your physician if the pain or fever persists for more than 5 days. Do not exceed the recommended dose unless advised by a physician.

If abdominal pain, heartburn, nausea or vomiting, bloating, diarrhea or constipation, ringing or buzzing in the ears, nervousness, sleeplessness, dizziness, any change in vision, fluid retention, itching, skin rashes or any other side effect or unexplained symptom develops while taking Advil, discontinue use immediately and contact a physician. In case of overdose, call a Poison Control Centre or a doctor at once even if there are no symptoms.

### Advil Liqui-Gels (200 mg) Non-medicinal Ingredients

FD&C green No. 3, gelatin, polyethylene glycol, polyvinyl acetate phthalate, potassium hydroxide, propylene glycol, purified water, sorbitan, sorbitol, titanium dioxide.

### Advil Extra Strength Liqui-Gels (400 mg) Non-medicinal Ingredients

gelatin, iron oxide, lecithin, medium chain triglycerides, polyethylene glycol, polyvinyl acetate phthalate, potassium hydroxide, propylene glycol, purified water, sorbitan, sorbitol.